


WHAT IS CLAIMED IS:

1. A compound having the formula:



(I)

wherein,

L^1 and L^2 are members independently selected from a bond, -O-, -S-, S(O)-, -S(O₂)-, -C(O)-, -C(O)O-, -C(O)NH-, substituted or unsubstituted alkylene, and substituted or unsubstituted heteroalkylene;

the dashed line b is optionally a bond;

the ring A is a member selected from substituted or unsubstituted 5 to 6 membered heterocycloalkyl, and substituted or unsubstituted heteroaryl;

R^1 is a member selected from hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, -OR^{1A}, -NR^{1C}R^{1D}, -C(O)NR^{1C}R^{1D}, -C(O)OR^{1A}, wherein

R^{1A} is a member selected from hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, and substituted or unsubstituted heteroaryl;

R^{1C} and R^{1D} are members independently selected from substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, and substituted or unsubstituted heteroaryl,

wherein R^{1C} and R^{1D} are optionally joined to form a substituted or unsubstituted ring with the nitrogen to which they are attached,

wherein said ring optionally comprises an additional ring
nitrogen, and

R^2 is a member selected from substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-S(O_2)R^{2A}$, $-S(O_2)NR^{2B}R^{2C}$, and $=NOR^{2D}$, wherein R^{2A} , R^{2B} , R^{2C} , and R^{2D} are members independently selected from substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, and substituted or unsubstituted heteroaryl.

2. The compound of claim 1, wherein A is a member selected from:
unsubstituted 5 to 6 membered heterocycloalkyl comprising at least one heteroatom selected from N, O and S;
substituted 5 to 6 membered heterocycloalkyl comprising 1 to 3 substituents and at least one ring heteroatom selected from N, O and S;
unsubstituted aryl comprising at least one heteroatom selected from N, O and S; and
substituted aryl comprising 1 to 3 substituents and at least one ring heteroatom selected from N, O and S.


3. The compound of claim 1, wherein A is a member selected from substituted or unsubstituted pyrrolidinyl, substituted or unsubstituted pyrrolyl, substituted or unsubstituted pyrazolyl, substituted or unsubstituted imidazolyl, substituted or unsubstituted furanyl, substituted or unsubstituted oxazolyl, substituted or unsubstituted isoxazolyl, substituted or unsubstituted thienyl, substituted or unsubstituted thiazolyl, substituted or unsubstituted isothiazolyl, substituted or unsubstituted pyridinyl, substituted or unsubstituted pyrimidinyl, and substituted or unsubstituted pyrazinyl.

4. The compound of claim 1, wherein A is a substituted or unsubstituted pyrazolyl.

5. The compound of claim 1, wherein A is substituted with a member selected from hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroaryl, substituted or unsubstituted aryl, $-NR^{3A}R^{3B}$, and $-OR^{3C}$, wherein R^{3A} and R^{3B} are members independently selected from hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted heterocycloalkyl, and substituted or unsubstituted heteroaryl, wherein R^{3A} and R^{3B} are optionally joined to form a substituted or unsubstituted ring with the nitrogen to which they are attached, wherein said ring optionally comprises an additional ring heteroatom, and R^{3C} is a member selected from substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, and substituted or unsubstituted heteroaryl.

6. The compound of claim 5, wherein A is substituted with a member selected from hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted aryl, and substituted or unsubstituted heteroaryl.

7. The compound of claim 1 having the formula



wherein,

the dashed ring represents unsaturated, partially saturated, or fully saturated bonds within ring E;

Z^1 is a member selected from $-NR^5-$, $=N-$, $-O-$, and $-S-$, wherein

R^5 is a member selected from hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted

heterocycloalkyl, substituted or unsubstituted heteroaryl, and substituted or unsubstituted aryl;

Z^2 is a member selected from $-\text{CR}^{6A}\text{R}^{6B}-$, $=\text{CR}^{6A}-$, $-\text{C}(\text{O})-$, $-\text{NR}^{6C}-$, $=\text{N}-$, $-\text{O}-$, $-\text{S}-$, $-\text{CR}^{6A}\text{R}^{6B}-\text{NR}^{6C}-$, $=\text{CR}^{6A}-\text{NR}^{6C}-$, $-\text{CR}^{6A}=\text{N}-$, $-\text{CR}^{6A}\text{R}^{6B}-\text{N}=\text{}$, and $=\text{CR}^{6A}-\text{N}=\text{}$, wherein

R^{6C} is a member selected from hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, and substituted or unsubstituted heteroaryl,

R^{6A} and R^{6B} are members independently selected from hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroaryl, substituted or unsubstituted aryl, $-\text{NR}^{6A1}\text{R}^{6A2}$, and $-\text{OR}^{6A3}$, wherein

R^{6A1} and R^{6A2} are members independently selected from hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, and substituted or unsubstituted heteroaryl, wherein

R^{6A1} and R^{6A2} are optionally joined to form a substituted or unsubstituted ring with the nitrogen to which they are attached, wherein said ring optionally comprises an additional ring heteroatom, and

R^{6A3} is a member selected from substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, and substituted or unsubstituted heteroaryl,

wherein R^{6A} and R^{6C} are optionally joined together to form a substituted or unsubstituted ring, wherein said ring optionally comprises an additional ring heteroatom;

Z^3 is a member selected from $-\text{CR}^{7A}\text{R}^{7B}-$, $=\text{CR}^{7A}-$, $-\text{C}(\text{O})-$, $-\text{NR}^{7C}-$, $=\text{N}-$, $-\text{O}-$, and $-\text{S}-$, wherein

R^{7C} is a member selected from hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroaryl, and substituted or unsubstituted aryl,

R^{7A} and R^{7B} are independently selected from hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroaryl, substituted or unsubstituted aryl, $-NR^{7A1}R^{7A2}$, and $-OR^{7A3}$, wherein R^{7A1} and R^{7A2} are members independently selected from hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, and substituted or unsubstituted heteroaryl, wherein R^{7A1} and R^{7A2} are optionally joined to form a substituted or unsubstituted ring with the nitrogen to which they are attached, wherein said ring optionally comprises an additional ring heteroatom, and

R^{7A3} is a member selected from substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, and substituted or unsubstituted heteroaryl;

wherein R^5 is optionally joined with R^{6A} or R^{6C} to form a substituted or unsubstituted ring, wherein said ring optionally comprises an additional ring heteroatom;

wherein R^{7A} is optionally joined with R^{6A} or R^{6C} to form a substituted or unsubstituted ring, wherein said ring optionally comprises an additional ring heteroatom; and

wherein R^{7C} is optionally joined with R^{6A} or R^{6C} to form a substituted or unsubstituted ring, wherein said ring optionally comprises an additional ring heteroatom.

8. The compound of claim 7, wherein

Z^1 is $-NR^5-$;

Z^2 is $=N-$; and


Z^3 is $=CR^{7A}-$.

1 9. The compound of claim 8, wherein

2 R^{7A} is hydrogen; and

3 R^5 is a member selected from hydrogensubstituted or unsubstituted aryl,
4 substituted or unsubstituted heteroaryl, substituted or unsubstituted arylalkyl and substituted
5 or unsubstituted heteroarylalkyl.

1 10. The compound of claim 7, wherein R^5 has the formula:



(VI)

3 wherein,

4 R^{5A} is a member selected from hydrogen, halogen, $-OR^{5A1}$, $-NR^{5A2}R^{5A3}$,
5 $-S(O_2)NR^{5A2}R^{5A3}$, $-CN$, substituted or unsubstituted alkyl, substituted or
6 unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl,
7 substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted
8 aryl, and substituted or unsubstituted heteroaryl, wherein

9 R^{5A1} is a member selected from hydrogen, substituted or unsubstituted
10 alkyl, substituted or unsubstituted heteroalkyl, substituted or
11 unsubstituted cycloalkyl, substituted or unsubstituted
12 heterocycloalkyl, substituted or unsubstituted aryl, and substituted or
13 unsubstituted heteroaryl, and

14 R^{5A2} and R^{5A3} are members independently selected from hydrogen,
15 substituted or unsubstituted alkyl, substituted or unsubstituted
16 heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or
17 unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, and
18 substituted or unsubstituted heteroaryl;

19 m is an integer from 0 to 10; and

20 n is an integer from 1 to 5.

1 11. The compound of claim 10, wherein

2 n is 1;

3 m is 0 or 1; and

4 R^{5A1} , R^{5A2} and R^{5A3} are hydrogen.

1 12. The compound of claim 7, wherein


Z^1 is $-NR^5-$;
 Z^2 is $=CR^{6A}-$; and
 Z^3 is $=N-$.

13. The compound of claim 12, wherein R^5 is a member selected from hydrogen and substituted or unsubstituted aryl.

14. The compound of claim 8, wherein R^5 and R^{7A} are hydrogen and b is a bond.

15. The compound of claim 1, wherein R^1 is a member selected from substituted or unsubstituted (C_1-C_{10}) alkyl, substituted or unsubstituted 2-10 membered heteroalkyl, substituted or unsubstituted (C_3-C_7) cycloalkyl, substituted or unsubstituted 3-7 membered heterocycloalkyl, substituted or unsubstituted aryl, and substituted or unsubstituted heteroaryl.

16. The compound of claim 1, wherein R^1 has the formula:



(III)

wherein,

q is an integer selected from 1 to 5;

R^{1B} is a member selected from hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-NR^{1B1}R^{1B2}$, $-OR^{1B3}$, and $-C(O)NR^{1B4}R^{1B5}$ wherein

R^{1B1} and R^{1B2} are members independently selected from hydrogen, substituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted heterocycloalkyl, and substituted or unsubstituted heteroaryl, wherein R^{1B1} and R^{1B2} are optionally joined to form a substituted or unsubstituted ring with the nitrogen to which they are attached, wherein said ring optionally comprises an additional ring heteroatom, and

R^{1B3} is a member selected from hydrogen,


substituted or unsubstituted heteroalkyl comprising a nitrogen,
 substituted or unsubstituted heterocycloalkyl comprising a ring
 nitrogen,
 substituted or unsubstituted heteroaryl comprising a ring nitrogen,
 and
 alkyl substituted with a substituted or unsubstituted heteroalkyl
 comprising a nitrogen, substituted or unsubstituted
 heterocycloalkyl comprising a ring nitrogen, and substituted or
 unsubstituted heteroaryl comprising a ring nitrogen; and
 R^{1B4} and R^{1B5} are members independently selected from
 hydrogen,
 substituted or unsubstituted heteroalkyl comprising a nitrogen,
 substituted or unsubstituted heterocycloalkyl comprising a ring nitrogen,
 substituted or unsubstituted heteroaryl comprising a ring nitrogen, and
 alkyl substituted with a substituted or unsubstituted heteroalkyl
 comprising a nitrogen, substituted or unsubstituted heterocycloalkyl
 comprising a ring nitrogen, and substituted or unsubstituted
 heteroaryl comprising a ring nitrogen, wherein
 R^{1B4} and R^{1B5} are optionally joined to form a substituted or
 unsubstituted ring with the nitrogen to which they are attached,
 wherein said ring optionally comprises a heteroatom.

17. The compound of claim 16, wherein

q is an integer selected from 1 to 3;

R^{1B} is a member selected from hydrogen, substituted alkyl, substituted or
 unsubstituted heteroalkyl, substituted cycloalkyl, substituted or
 unsubstituted heterocycloalkyl, substituted aryl, and substituted or
 unsubstituted heteroaryl.

18. The compound of claim 16, wherein R^1 has the formula:



(IV)

wherein,

R^{1B} is a member selected from hydrogen, $-NR^{1B1}R^{1B2}$, $-OR^{1B3}$, substituted or
 unsubstituted (C_1 - C_{10}) alkyl, substituted or unsubstituted 2-10

6 membered heteroalkyl, substituted or unsubstituted (C₃-C₇)cycloalkyl,
7 substituted or unsubstituted 3-7 membered heterocycloalkyl, substituted
8 or unsubstituted aryl, and substituted or unsubstituted heteroaryl.

1 **19.** The compound of claim 16, wherein R^{1B} is a member selected from
2 -C(O)NR^{1B4}R^{1B5} and substituted or unsubstituted heteroaryl comprising a ring nitrogen,
3 wherein

4 R^{1B4} and R^{1B5} are members independently selected from
5 hydrogen,
6 substituted or unsubstituted heteroalkyl comprising a nitrogen,
7 substituted or unsubstituted heterocycloalkyl comprising a ring nitrogen,
8 substituted or unsubstituted heteroaryl comprising a ring nitrogen, and
9 alkyl substituted with a substituted or unsubstituted heteroalkyl
10 comprising a nitrogen, substituted or unsubstituted heterocycloalkyl
11 comprising a ring nitrogen, and substituted or unsubstituted
12 heteroaryl comprising a ring nitrogen, wherein
13 R^{1B4} and R^{1B5} are optionally joined to form a substituted or
14 unsubstituted ring with the nitrogen to which they are attached,
15 wherein said ring optionally comprises a heteroatom.


1 **20.** The compound of claim 19, wherein R^{1B1}, R^{1B2}, R^{1B3}, R^{1B4} and
2 R^{1B5} are members independently selected from hydrogen and a substituted or unsubstituted
3 ring, wherein said ring optionally comprises a nitrogen atom and at least one additional ring
4 heteroatom.

1 **21.** The compound of claim 1, wherein R² is a member selected from
2 substituted or unsubstituted (C₁-C₁₀) alkyl, substituted or unsubstituted 2-10 membered
3 heteroalkyl, substituted or unsubstituted (C₃-C₇) cycloalkyl, substituted or unsubstituted 3-7
4 membered heterocycloalkyl, substituted or unsubstituted aryl, and substituted or
5 unsubstituted heteroaryl.

1 **22.** The compound of claim 1, R^{2A}, R^{2B}, R^{2C}, and R^{2D} are members
2 independently selected from substituted or unsubstituted (C₁-C₁₀) alkyl, substituted or
3 unsubstituted 2-10 membered heteroalkyl, substituted or unsubstituted (C₃-C₇) cycloalkyl,

substituted or unsubstituted 3-7 membered heterocycloalkyl, substituted or unsubstituted aryl, and substituted or unsubstituted heteroaryl.

23. The compound of claim 1, R^2 has the formula:



(V)

wherein,

R^{2G} is a member selected from hydrogen, halogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, and substituted or unsubstituted heteroaryl;

J is a substituted or unsubstituted ring selected from substituted or unsubstituted (C_3 - C_7) cycloalkyl, substituted or unsubstituted 3-7 membered heterocycloalkyl, substituted or unsubstituted aryl, and substituted or unsubstituted heteroaryl;

t is an integer from 0 to 5; and

X is a member selected from a bond, $-S(O_2)-$, and $-S(O_2)N^{2I}-$, wherein

R^{2I} is a member selected from hydrogen, substituted or unsubstituted alkyl, and substituted or unsubstituted heteroalkyl.

24. The compound of claim 23, wherein

R^{2G} is a member selected from hydrogen, substituted or unsubstituted (C_1 - C_{10}) alkyl, substituted or unsubstituted 2-10 membered heteroalkyl, substituted or unsubstituted (C_3 - C_7) cycloalkyl, substituted or unsubstituted 3-7 membered heterocycloalkyl, substituted or unsubstituted aryl, and substituted or unsubstituted heteroaryl;

J is a substituted or unsubstituted ring selected from substituted or unsubstituted 3-7 membered heterocycloalkyl, substituted or unsubstituted aryl, and substituted or unsubstituted heteroaryl;

t is 1; and


R^{2I} is hydrogen.

25. The compound of claim 23, wherein R^{2G} is a branched or unbranched (C_1-C_{10}) alkyl.

26. The compound of claim 23, wherein X is $-S(O_2)-$.

27. The compound of claim 1, wherein L^1 and L^2 are members independently selected from a bond and unsubstituted (C_1-C_6) alkylene.

28. The compound of claim 1, wherein
the dashed line b is a bond;
 R^1 is substituted or unsubstituted benzyl; and
 R^2 has the formula:



(V)

wherein,

R^{2G} is a member selected from hydrogen, halogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, and substituted or unsubstituted heteroaryl,

J is a substituted or unsubstituted ring selected from substituted or unsubstituted (C_3-C_7) cycloalkyl, substituted or unsubstituted 3-7 membered heterocycloalkyl, substituted or unsubstituted aryl, and substituted or unsubstituted heteroaryl,

t is an integer from 0 to 5, and

X is $-S(O_2)-$;

L^1 is a bond; and

L^2 is a bond.

29. A method of treating a disorder or condition through modulating a glucocorticoid receptor, the method comprising administering to a subject in need of such treatment, an effective amount of the compound of one of claims 1-28.

1 **30.** A method of treating a disorder or condition through antagonizing a
2 glucocorticoid receptor, the method comprising administering to a subject in need of such
3 treatment, an effective amount of the compound of one of claims **1-28**.

1 **31.** A method of modulating a glucocorticoid receptor including the steps
2 of contacting a glucocorticoid receptor with an effective amount of the compound of one of
3 claims **1-28** and detecting a change in the activity of the glucocorticoid receptor.

1 **32.** A pharmaceutical composition comprising a pharmaceutically
2 acceptable excipient and the compound of one of claims **1-28**.